



Examples 8.7 – Modeling Accumulated Change with the TI-84

Consider the function $f(x) = (x-2)^3 - 5x + 12$ for x in the interval $[0, 5]$. In the function menu, enter $Y_1 = (X-2)^3 - 5X + 12$, and set the viewing window as $[0, 5] \times [-6, 7]$.

If we want the net area bounded by the graph of f and the x -axis on an interval, then we use option 7 in the CALCULATE menu.

1: To find $\int_1^3 f(x) dx$, for instance, choose 7: $\int f(x) dx$, input $x = 1$ and press **[ENTER]**, and then input $x = 3$ and press **[ENTER]**. The region bounded by the graph and the x -axis on the chosen interval will be shaded. The result will be the net area.

If we want the total area, then we must work with the absolute value of f . In the function menu, enter $Y_2 = \text{abs}(Y_1(X))$, and set the viewing window as $[0, 5] \times [-6, 7]$. To get the absolute value function, press **[2ND] [0] [ENTER]** or **[MATH] [►] [ENTER]**. To get Y_1 , press **[VARS] [►] [ENTER] [ENTER]**. Now we use option 7 in the calculate menu as before.

7: To find $\int_1^3 |f(x)| dx$, for instance, choose 7: $\int f(x) dx$, toggle to Y_2 , input $x = 1$ and press **[ENTER]**, and then input $x = 3$ and press **[ENTER]**. The region bounded by the graph and the x -axis on the chosen interval will be shaded. The result will be the total area.